

Effect of herbicide (sodium penta chloro phanate) on pollen germination and tube length of stored pollen of Apocynaceae: Further evidence of a criticism of Banerji and Gangulee (1937), Brewbaker and Kwack (1963), Sudhakaran (1967- Ph.D.Thesis), Dharurkar (1971- Ph.D. Thesis), Nair, Nambudiri and Thomas (1973), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980- Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil and Haldar (1980), Shetye (1982- Ph.D. Thesis) and Giridhar (1984- Ph.D. Thesis)—A critical review

# S.A. Salgare

Received on: 15-09-2008

# Accepted on: 15-01-2009 Abstract

Even the lowest concentration (10<sup>-17</sup>mg/ml) of sodium penta chloro phanate tried suppressed the germination of pollen of F and F-24 series of red-flowered cultivar of *Nerium odorum* and F-48 and F-72 series of pink-flowered cultivar of *Catharanthus roseus*. The herbicide stimulated the germinataion of pollen of successive flowers of all the cultivars of the Apocynaceae throughout the experiment. However, it stimulated the tube growth of only 3 out of 10 series.

Keywords:- Palynology, Toxicology, Environmental Sciences, Herbicides

#### Introduction

Herbicides drastically reduced pollen germination as well as tube growth. It was therefore important to study the effect of such chemicals on germination as well as tube growth since inhibitory effects of these chemicals eventually reduce fruit and seed-set.

#### Materials and Method

Pollen of successive flowers (viz. F, F-24, F-48, F-72 series i.e. open flowers and the flower buds which require 24, 48, 72 hours to open respectively) of 5 cultivars of Apocynaceae e.g. red-, pink- and white-flowered cultivars of Nerium odorum Soland. and pink- and white-flowered cultivars of Catharanthus roseus (L.) G. Don. were collected soon after the

#### **Author Address**

Salgare Research Foundation Pvt. Ltd., Prathamesh Society, Shivaji Chowk, Karjat

dehiscence of anthers in the open flowers and stored at room temperature (22-31.8°C) having RH 57% and in diffuse laboratory light at the Department of Botany, Govt. Institute of Science, Mumbai. Germination of stored pollen grains of successive flowers was made soon after the dehiscence of anthers and with 2 hours intervals for the first 10 hours in the optimum concentrations of sucrose (acts as control) as well as in the optimum concentrations of sucrose supplemented with the optimum concentrations of sodium penta chloro phanate (Table 1). Observations were recorded 24 hours after incubation. For each experiment a random count of 200 grains was made to determine the percentage of pollen germination. For measurement of length of pollen tubes, 50 tubes were selected randomly and measured at a magnification of 100x.

Table 1: Effect of sodium penta chloro phenate on stored pollen on their germination and tube length of

| five cult | tivars of | Apocynaceae |
|-----------|-----------|-------------|
|-----------|-----------|-------------|

| Cultivars                | SF   | PV | SC | нс    | PG & TLSADA |    |          |      | TRADAFMPGTAL |    |   |      |    |    |    |      |
|--------------------------|------|----|----|-------|-------------|----|----------|------|--------------|----|---|------|----|----|----|------|
|                          |      |    |    |       | %PG         |    | TL in mm |      | С            |    |   |      | T  |    |    |      |
|                          |      |    |    |       | C           | T  | С        | T    | H            | PG | H | TL   | H  | PG | H  | TL   |
| N. odorum pink-flowered  | F    | 80 | 50 | 10-17 | 35          | 37 | 1485     | 745  | 8            | 52 | 0 | 1485 | 4  | 45 | 4  | 836  |
| N. odorum red-flowered   | F    | 74 | 20 | Ng    | 20          | Ng | 1250     | Ng   | 6            | 36 | 0 | 1250 | Ng | Ng | Ng | Ng   |
| N. odorum white-flowered | F    | 62 | 50 | 10-17 | 20          | 40 | 675      | 148  | 4            | 24 | 0 | 728  | 2  | 45 | 4  | 210  |
| C. roseus pink-flowered  | F    | 90 | 20 | 10-17 | 60          | 74 | 1575     | 1800 | 6            | 72 | 0 | 1575 | 0  | 74 | 0  | 1800 |
| C. roseus white-flowered | F    | 88 | 20 | 10-17 | 40          | 64 | 1256     | 1220 | 8            | 75 | 0 | 1256 | 4  | 75 | 0  | 1220 |
| N. odorum red-flowered   | F-24 | 74 | 20 | Ng    | 06          | Ng | 485      | Ng   | 6            | 8  | 0 | 485  | Ng | Ng | Ng | Ng   |
| C. roseus pink-flowered  | F-24 | 90 | 50 | 10-17 | 28          | 29 | 240      | 425  | 6            | 36 | 4 | 300  | 0  | 29 | 0  | 425  |
| C. roseus white-flowered | F-24 | 88 | 50 | 10-17 | 16          | 56 | 248      | 786  | 6            | 50 | 6 | 264  | 2  | 60 | 0  | 789  |
| C. roseus pink-flowered  | F-48 | 90 | 50 | Ng    | 14          | Ng | 95       | Ng   | 4            | 16 | 0 | 95   | Ng | Ng | Ng | Ng   |
| C. roseus pink-flowered  | F-72 | 90 | 80 | Ng    | 10          | Ng | 65       | Ng   | 0            | 10 | 0 | 65   | Ng | Ng | Ng | Ng   |

C, in control; HC, herbicide concentrations in mg/ml; Ng, no germination of pollen even 24 hours of their sowing; PG, pollen germination; PG&TLSADA, Pollen germination & tube length in the sets which were set soon after dehiscence of anthers; PV, pollen viability in %; SC, sucrose concentrations in %; SF, successive flowers; TL, tube length; TRADAFMPGTL, Time required after dehiscence of anthers for maximum pollen germination & tube length.

## Results and Discussion

Pollen viability is a subject that has a great deal of practical as well as theoretical interest. In the present investigation even the different cultivars of the same species showed the variations in the percentage of their pollen viability (Table 1). Reduced pollen viability has been interpreted as an indication of suspected hybridity in wild populations. Nevertheless, variations in pollen viability may affect the breeding systems of the species concerned, and if the pollen viability can be altered by the environment, then the breeding system itself may be under some degree of environmental control.

Potentiality of pollen germinability was recorded in F series of all the 5 cultivars of the Apocynaceae studied. It was the pollen of F-24 series of red-flowered cultivar of *Nerium odorum* and both the cultivars of *Catharanthus roseus* found germinated in the optimum concentrations of sucrose. It should be pointed out that the pollen of F-48 and F-72 series of pink-flowered cultivar of *C. roseus* showed their germination in the optimum concentrations of

sucrose. Thus the potentiality of pollen germinability in Apocynaceae was observed in 10 out of 20 series investigated (Table 1).

As a rule the percentage of pollen germination is always less than the pollen viability (Table 1). However, Banerji and Gangulee (1937) and Dharurkar (1971) reported higher percentage of pollen germination than the pollen viability in *Eichhornia crassipes*. The claim of Banerji and Gangulee (1937) and Dharurkar (1971) is challenged by Salgare (1986c, 1995, 2000b, 2006a, 2006j, 2006m, 2006q, 2006s, 2007b-2007c, 2007e, 2007f, 2007g, 2007i and 2007j) who stated that the observations of Banerji and Gangulee (1937) and Dharurkar (1971) are exaggerating.

Germination of pollen of F-72 series of pink-flowered cultivar of Catharanthus roseus in vitro culture of sucrose was noted in the present investigation. However, Palathingal (1990) failed to germinate the pollen of F-72 series of pink-flowered cultivar of C. roseus in Brewbaker and Kwack's (1963) culture medium. This proves that the culture medium is also having the bearing on the germination of pollen.



This also confirms that Brewbaker and Kwack's (1963) culture medium is not ideal for pollen cultures. This was also pointed out earlier by Salgare (2006a, 2006j, 2006g, 2006s, 2006u, 2007e, 2007f and 2007h). Even the lowest concentration (10-17 mg/ml) of sodium penta chloro phanate tried suppressed the germination of pollen of F and F-24 series of redflowered cultivar of Nerium odorum and F-48 and F-72 series of pink-flowered cultivar of Catharanthus roseus (Table 1). Sharma (1984) stated that even the lowest concentration (10-17 mg/ ml) of sodium penta chloro phanate tried prevented the germination of pollen of F series of white cascade, sonata, F-24 series of white cascade, duet, sonata and F-48 series of red as well as white cascade. All these are the cultivars of Petunia grandiflora. This proves that the pollen of the said series are highly sensitive and acts as an ideal indicators of pollution. Thus it is confirmed that the pollen development and activity are more sensitive indicators of adverse factors in the botanical environment and the use of an entire vascular plant (Berg, 1973; Brandt, 1974; Vick and Bevan, 1976; Rasmussan, 1977; Navara et al. 1978; Mhatre, 1980; Mhatre et al. 1980; Shetye, 1982 and Giridhar, 1984) as an indicator of pollution is a very crude method and rather a wrong choice. There is no evidence of any entire vascular plant exhibiting this much degree of sensitivity. This is confirmed in the present critical review (Table 1) as well as by the previous extensive work of Salgare (1983, 1984,1985a, 1985b, 1985c, 1986a, 1986e, 1986f, 2000a, 2001a, 2001b, 2005b, 2005d, 2005e, 2006a, 2006f, 2006j, 2006l, 2006o, 2006q, 2006u, 2007b, 2007c, 2007d, 2007e, 2007f, 2007g, 2007h, 2007i, 2007i, 2007k, 2007l), Salgare and Theresa Sebastian (1986), Salgare and Phunguskar (2002), Salgare and Singh (2002, 2006a, 2006b), Salgare and Pathak (2005) and Sharma (1984).

Sodium penta chloro phanate stimulated the germination of pollen of successive flowers of all the cultivars of the apocynaceae throughout the

experiment. However, it stimulated the tube growth of only 3 out of 10 series (Table 1). Maximum pollen germination was noted in all the 10 series except for F-72 series of pink-flowered cultivar of Catharanthus roseus with the stored pollen in vitro culture of sucrose. Pollen of F-72 series of pink-flowered cultivar of C. roseus showed an equal percentage of germination of pollen with that stored pollen. The time interval of the period of storage ranges right from 4 to 8 hours. However, the pollen of F-72 series of pink-flowered cultivar of C. roseus showed the highest germination in the sets which were set soon after the dehiscence of the anthers (Table 1). The herbicide stimulated the germination of stored pollen in 4 out of 10 series, while the tube growth was stimulated in 2 out of 10 series of the Apocynaceae. In vitro culture of sucrose the longest pollen tubes were noted in 5 out of 10 series in the sets which were set soon after the dehiscence of the anthers. In vitro culture of sucrose supplemented with the herbicide the maximum germination of pollen was noted in 6 out of 7 series in the sets which were sets soon after the dehiscence of the anthers. However, the longest pollen tubes were noted in 3 out of 7 series in the sets set soon after the dehiscence of anther (Table 1). Thus it is confirmed that the pollen germination and tube elongation are two distinct processes. However, Nair et al. (1973) stated that the pollen germination and tube elongation is one and the same process. Present work (Table 1) as well as previous extensive work of Salgare (1979, 1983, 1986d, 2004, 05a, 2005c, 2006e, 2006j, 2006k, 200l, 2006s, 2007i, 2007j), Salgare and Bindu (2002, 2005) and Salgare and Tessy Mol Antony (2005a, 2005b) it could be concluded that the observations of Nair et al. (1973) are superficial and misleading.

In this connection it should be pointed out that Sudhakaran (1967) stated that in *Vinca rosea* L. {Catharanthus roseus (L.) G. Don.} besides pollen grains which produced single pollen tube, it has also been noticed that tetraploid grains frequently produce more than one pollen tube. Pollen tubes



are branched quite frequently. Aberrations of this type in the pollen tube development are not observed in diploid pollen tubes, but quite frequently met with the pollen grains of irradiated plants. Salgare (1983) made it very clear that Sudhakaran (1967) had failed to trace out the branched pollen tubes and polysiphonous condition which is fairly common even in diploid pollen grains. Apart from this Sudhakaran (1967) was not able to report the various types of pollen tube deformities either with diploid or tetraploid grains. Present investigation as well as the extensive work of Salgare (1983, 1986b, 2006b, 2006c, 2006d, 2006h, 2006j, 2006n, 2006p, 2006g, 2006s, 2006t, 2007a, 2007b, 2007c, 2007e, 2007f, 2007h, 2007i) also proved that Sudhakaran's (1967) observations are superficial and misleading.

### References

- Banerji, I. and Gangulee, H.C., 1937. Spermatogenesis in Eichhornia crassipes Solms. J. Indian Bot. Soc., 16: 289-296.
- Berg, H., 1973. Plants as indicators of air pollution. *Toxicol.*, 1: 79-89.
- Brandt, C.C., 1974. *Indicators of environmental quality*, In: W.A. Thomas (ed.) Plenum Press, New York. pp: 101.
- Brewbaker, J.L. and Kwack, B.H., 1963. The essential role of Ca ion in pollen germination and pollen tube growth. *Amer. J. Bot.*, '50: 859-865.
- Dharurkar, R.D., 1971. Effects of herbicides on the cytomorphology of *Eichhornia crassipes* Solms (Mart.). *Ph.D. Thesis, Univ. Bombay.*
- Giridhar, B.A., 1984. Study of interactions between industrial air pollutants and plants. Ph.D. Thesis, Univ. Bombay.
- Mhatre, G.N., 1980. Studies in responses to heavy metals in industrial Environment. Ph.D. Thesis, Univ. Bombay.
- Mhatre, G.N., Chaphekar, S.B., Ramani Rao, I.V., Patil, M.R. and Haldar, B.C., 1980. Effect of industrial pollution on the Kalu river ecosystem. *Environ. Pollut. Series A*, 23: 67-78.
- Nair, P.K.K., Nambudiri, E.M.V. and Thomas, M.K., 1973.

  A note on pollen germination at various stages of

- development of flower buds of Balsam (Impatians balsam). J. Palyno., 4: 29-33.
- Navara, J., Horvath, I. and Kaleta, M., 1978. Contribution to the determination of limiting of values of SO<sub>2</sub> for vegetation in the region of Bratislava. *Environ. Pollut.* Series A, 16: 249-262.
- Rasmussan, L., 1977. Epiphytic bryophytes as indicators of changes in the background levels of air borne metals from 1951 to 1975. Environ. Pollut. Series A, 14: 34-45.
- Salgare, S.A., 1979. A Criticism of Nair, Nambudiri and Thomas' (1973) paper entitled, 'A note on pollen germination at various stages of development of flower buds of balsam (*Impatiens balsam*)'. Proc. of 66th Session of the Indian Sci. Cong. held on January 3-7, 1979 at Hyderabad, Section of Bot., 3:61-62, Abstract No. 131.
- Salgare, S.A., 1983. Pollen physiology of Angiosperms I. Ph. D. Thesis, Univ. Bombay.
- Salgare, S.A., 1984. Further Evidence of a Criticism of the Hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis). Proc. 4th Indian Palyno. Conf., held on March 19-21, 1984 at the Deptt. of Environ. Sci., Andhra Univ., Visakhapatnam 530 003, Abstract No. S III-06.
- Salgare, S.A., 1985a. A Criticism on the Findings of Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis). Proc. 2<sup>nd</sup> Ann. Conf. of the Nat. Environ. Sci. Acad., held on May 25-27, 1985 at the Awadh Univ., Faizabad. Abstract No. 2.
- Salgare, S.A., 1985b. Further Evidence of the Findings of Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis).
  Proc. Nat. Symp. on Environ. Sci. and Warm Water Aquaculture Finfish, held on November 6-9, 1985 at the Deptt. of Zool., M.J.K. College of Bihar Univ., Bettiah 845 438, Abstract No. 2.
- Salgare, S.A., 1985c. A Criticism on the Findings of Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Pattil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis). Proc. Nat. Symp. on Environ. Sci. and Warm Water Aquaculture Finfish, held on November 6-9, 1985



- at the Deptt. of Zool., M.J.K. College of Bihar Univ., Bettiah 845 438, Abstract No. 73.
- Salgare, S.A., 1986a. Effect of herbicides on pollen physiology of Petunia axillaris BSP. Ph.D. Thesis, World Univ.
- Salgare, S.A., 1986b. A Criticism on Ph.D. Thesis of Sudhakaran (1967) entitled, 'Cytogenetic studies in Vinca rosea Linn.'. Proc. 1st Nat. Symp. on Laviron. Biol., held on December 30-31, 1986 at Deptt. Zool. and Microbiol., S. K. Univ., Anantapur – 515 003, Abstract No. 5.
- Salgare, S.A., 1986c. A Criticism on Ph.D.Thesis of Dharurkar (1971) entitled, 'Effect of herbicides on the cytomorphology of *Eichhornia crassipes* Solms (Mart.). Ph.D. Thesis, Univ. of Bombay. Proc. 1st Nat. Symp. on Environ. Biol., held on December 30-31, 1986 at Deptt. Zool. and Microbiol., S. K. Univ., Anantapur 515 003, Abstract No. 6.
- Salgare, S.A., 1986d. A Criticism on the Findings of Nair, Nambudiri and Thomas (1973). Proc. 1st Nat. Symp. on Environ. Biol., held on December 30-31, 1986 at Deptt. Zool. and Microbiol., S. K. Univ., Anantapur – 515 003, Abstract No. 13.
- Salgare, S.A., 1986e. A Criticism on Ph.D. Thesis of Giridhar (1984) entitled, 'Study of Interactions between industrial air Pollutants and plants'. Ph.D.Thesis, Univ. Bombay. Proc. 1st Nat. Symp. on Environ. Biol., held on December 30-31, 1986 at Deptt. Zool. and Microbiol., S. K. Univ., Anantapur-515 003, Abstract No. 14.
- Salgare, S.A., 1986f. A Criticism on Ph.D. Thesis of Shetye (1982) entitled, 'Effect of Heavy Metals on plants'. Ph.D. Thesis, Univ. Bombay. Proc. 1st Nat. Symp. on Environ. Biol., held on December 30-31, 1986 at Deptt. Zool. and Microbiol., S.K. Univ., Anantapur– 515 003, Abstract No. 16.
- Salgare, S.A., 1995. Interesting observations on the physiology of pollen of Eichhornia crassipes. Solam. (Mart). Proc.13th Nat. Symp. on Life Sci., Proceeded with Ann. Session of Indian Soc. of Life Sci., held on December 30-31, 1995. January 1, 1996 at CH. Charan Singh Univ., Meerut- 250 004, Abstract No. 83.
- Salgare, S.A., 2000a. Additional Evidence of a Criticism on the Hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis). pp 64-65, Proc. 1st Nat. Conf. on Recent Trends in Life Management,

- held on October 22-23, 2000 at Bipin Bihari P. G. College, Jhansi- 284 001 (U.P.). Abstract No. 92.
- Salgare, S.A., 2000b. A Criticism on the Findings of Banerji and Gangulee (1937) and Dharurkar (1971- Ph.D. Thesis). *Him. J. Env. Zool.*, 14: 159-160.
- Salgare, S.A., 2001a. Monitoring of herbicide (nitrofen) toxicity by using pollen as indicators – A Critical Review– I. Website www.microbiologyou.com. U.S.A's Publications (35 pages).
- Salgare, S.A., 2001b. Monitors of pollution A Criticism on the Hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldara (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review. V. Biojournal, 13: 39-44.
- Salgare, S.A., 2004. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose + 2,4-D) by some species of Apocynaceae – A Criticism to the Hypothesis of Nair, Nambudiri and Thomas (1973) – A Critical Review – IV. Him. J. Env. Zool., 18: 53-56.
- Salgare, S.A., 2005a. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose+2,4,5-T) by some cultivars of Apocynaceae and Further Evidence of A Criticism to the Hypothesis of Nair, Nambudiri and Thomas (1973) - A Critical Review. Flora and Fauna, 11: 42-44.
- Salgare, S.A., 2005b. Monitoring of herbicide (simazine) toxicity— A Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984- Ph.D. Thesis)—A Critical Review. Him. J. Env. Zool., 19: 69-71.
- Salgare, S.A., 2005c. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose + nitrofen) - A Criticism to the Hypothesis of Nair, Nambudiri and Thomas (1973)- A Critical Review. *Him. J. Env. Zool.*, 19: 93-95.
- Salgare, S.A., 2005d. Monitoring of herbicide (basalin EC) toxicity— A Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984-Ph.D.



- Thesis) A Critical Review. *Internat. J. Biosci. Reporter*, 3: 298-300.
- Salgare, S.A., 2005e. Monitoring of herbicide (dalapon) toxicity by using pollen as indicators Pollen of some cultivars of Apocynaceae and further evidence of a Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980- Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982- Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis)- A Critical Review. Flora and Fauna, 11: 49-50.
- Salgare, S.A., 2006a. Monitoring of herbicide (fernoxone) toxicity by using pollen as indicators—pollen of five cultivars of *Petunia axillaris*: Further evidence of a Criticism of Banerji and Gangulee (1937), Brewbaker and Kwack (1963), Dharurkar (1971- Ph.D. Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review. *Environ. Conservation J.*, 7: 1-7.
- Salgare, S.A., 2006b. Further evidence of a Criticism of the findings of Sudhakaran (1967-Ph.D.Thesis) and Katre and Ghatnekar (1978)\*. Internat. J. Biosci. Reporter, 4: 19-20.
- Salgare, S.A., 2006c. Effect of acrolein on pollen germination and tube growth of stored pollen of Apocynaceae and further evidence of a Criticism of the hypothesis of Sudhakaran (1967) - A Critical Review. Him. J. Env. Zool., 20: 137-139.
- Salgare, S.A., (2006d). Effect of fernoxone on pollen germination and tube growth of twelve hours stored pollen of Apocynaceae and further evidence of a Criticism of the hypothesis of Sudhakaran (1967). Asian J. Bio Sci., 1: 160.
- Salgare, S.A., 2006e. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose + sodium arsenite) and Further Evidence of a Criticism of the Hypothesis of Nair, Nambudiri and Thomas (1973) - A Critical Review. *Internat. J. Pland Sci.*, 1: 132-133.
- Salgare, S.A., 2006f. Monitoring of herbicide (basalin EC) toxicity by using pollen as indicators—Pollen of Petunlia axillaris and further evidence of a Criticism of the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil,

- Haldar (1980), Shetye (1982- Ph.D. Thesis) and Giridhar (1984- Ph.D. Thesis) A Critical Review. *Him. J. Envi. Zool.*, 20: 269-271.
- Salgare, S.A., 2006g. Alteration of resting period of pollen of Apocynaceae by herbicide (nitrofen): Further evidence of a Criticism of Sudhakaran (1967-Ph.D.Thesis), Saoji and Chitaley (1972), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D.Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D.Thesis) and Giridhar (1984 Ph.D. Thesis). Internat. J. Agri. Sci., 3: 239-243.
- Salgare, S.A., 2006h. Effect of herbicide (acrolein) on pollen germination and tube growth of twelve hours stored pollen of five cultivars of Apocynaceae: Further evidence of a Criticism of Sudhakaran (1967-Ph.D.Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984-Ph.D. Thesis) A Critical Review\*. J. Natcon, 18: 283-290.
- Salgare, S.A., 2006i. Monitoring of herbicide (2,4-D) toxicity by using pollen as indicators Pollen of Apocynaceae and further evidence of a Criticism of the Hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review. *Internat. J. Plant Sci.*, 1: 134- 136.
- Salgare, S.A., 2006k. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose + atrataf 50w) - A Criticism to the hypothesis of Nair, Nambudiri and Thomas (1973) - A Critical Review. Internat. J. Biosci. Reporter, 4: 145-146.
- Salgare, S.A., 2006l. Evaluation of MH as male gametocide on Phaseolus aureus and a new method of plant breeding and further evidence of a Criticism of the hypothesis of Nair, Nambudiri, Thomas (1973), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980- Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis). Internat. J. Plant Sci., 1: 352-356.
- Salgare, S.A., 2006m. Further Evidence of a Criticism of the Findings of Banerji and Gangulee (1937) and



- Dharurkar (1971)\*. Internat. J. Biosci. Reporter, 4: 169-170.
- Salgare, S.A., 2006n. Effect of gramoxone on pollen germination and tube growth of twelve hours stored pollen of Apocynaceae and further evidence of a Criticism of the hypothesis of Sudhakaran (1967) - A Critical Review- II. Plant Archives, 6: 389-390.
- Salgare, S.A., 2006o. Alteration of resting period of pollen of five cultivars of *Petunia axillaris* BSP. by atrataf 50W: Further evidence of a Criticism of Saoji and Chitaley (1972), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984-Ph.D. Thesis)—A Critical Review. *J. Natcon*, 18: 353-360.
- Salgare, S.A., 2006p. Effect of atrataf 50W on pollen germination and tube growth of twelve hours stored pollen of Apocynaceae and further evidence of a Criticism of Sudhakaran (1967)- A Critical Review – 1\*. Internat. J. Biosci. Reporter, 4: 204- 206.
- Salgare, S.A., 2006q. Monitoring of herbicide (2,4-dinitrophenol) toxicity by using pollen as indicators—Pollen of five cultivars of Apocynaceae: Further evidence of a Criticism of Banerji and Gangulee (1937), Brewbaker and Kwack (1963), Sudhakaran (1967-Ph.D.Thesis), Dharurkar (1971 Ph.D. Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review\*, Res. Hunt, 1: 1-6.
- Salgare, S.A., 2006r. Alteration of resting period of pollen of five cultivars of *Petunia axillaris* BSP. by atrataf 50W: Further evidence of a Criticism of Saoji and Chitaley (1972), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984-Ph.D. Thesis) A Critical Review. *J. Natcon*, 18: 357-364.
- Salgare, S.A., 2006s. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose+ simazine) by five cultivars of Apocynaceae: Further evidence of a Criticism of Banerji and Gangulee (1937), Brewbaker and Kwack (1963), Sudhakaran (1967-Ph.D. Thesis), Dharurkar (1971-Ph.D. Thesis), Nair, Nambudiri and Thomas (1973), Berg (1973), Brandt (1974), Vick and Bevan (1976),

- Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis)- A Critical Review\*. Res. Hunt, 1: 146-155.
- Salgare, S.A., 2006t. Effect of herbicide (acrolein) on pollen germination and tube growth of twelve hours stored pollen of five cultivars of Apocynaceae: Furthe evidence of a Criticism of Sudhakaran (1967-Ph.D.Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D.Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982- Ph.D.Thesis) and Giridhar (1984-Ph.D.Thesis)- A Critical Review\*. J. Natcon, 18: 361-368.
- Salgare, S.A., 2006u. Alteration of resting period of pollen of five cultivars of Petunia axillaris BSP. by Gramoxone: Further evidence of a Criticism of Brewbaker and Kwack (1963), Saoji and Chitaley (1972), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984-Ph.D. Thesis) A Critical Review. Environ. Conservation J., 7: 51-57.
- Salgare, S.A., 2007a. Effect of Herbicide (sodium arsenite) on Pollen germination and Tube growth of twelve hours stored pollen of five cultivars of Apocynaceae and Further Evidence of a Criticism of Sudhakaran (1967): A Critical Review. Him. J. Env. Zool., 21: 171-173.
- Salgare, S.A., 2007b. Effect of herbicide (basalin EC) on pollen germination and tube growth of twelve hours stored pollen of five cultivars of Apocynaceae: Further evidence of a criticism of Banerji and Gangulee (1937), Sudhakaran (1967-Ph.D.Thesis), Dharurkar (1971 Ph.D. Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984 -Ph.D. Thesis) A Critical Review\*. Environ. Conservation J., 8: 1-8.
- Salgare, S.A., 2007c. Effect of Herbicide (simazine) on pollen germination and tube growth of twelve hours Stored Pollen of five cultivars of Apocynaceae: Further Evidence of a Criticism of Banerji and Gangulee (1937), Sudhakaran (1967-Ph.D.Thesis), Dharurkar (1971 - Ph.D. Thesis), Berg (1973), Brandt



- (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review. *Internat. J. Agric. Sci.*, 3: 313-317.
- Salgare, S.A., 2007d. Effect of atrataf 50W on the Rate of Pollen germination and tube growth of successive flowers of five cultivars of Petunia axillaris BSP. and Further Evidence of a Criticism of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review. Him. J. Env. Zool., 21: 377-379.
- Salgare, S.A., 2007e. Monitoring of Herbicide (nitrofen)
  Toxicity by Using Pollen as Indicators—Pollen of five cultivars of Apocynaceae [Further Evidence of a Criticism of Banerji and Gangulee (1937), Brewbaker and Kwack (1963), Sudhakaran (1967-Ph.D. Thesis), Dharurkar (1971 Ph.D. Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil and Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis)]. Plant Archives, 7: 397-400.
- Salgare, S.A., 2007f. Effect of Herbicide (MH) on pollen germination and tube growth of twelve hours stored pollen of Apocynaceae: Further Evidence of a Criticism of Banerji and Gangulee (1937), Brewbaker and Kwack (1963), Sudhakaran (1967-Ph.D.Thesis), Dharurkar (1971 Ph.D. Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review\*. *Internat. J. Plant Sci.*, 2: 215-219.
- Salgare, S.A., 2007g. Monitoring of herbicide (MH) toxicity by using pollen as indicators- Pollen of five cultivars of *Petunlia axillaris* BSP.: Further evidence of a criticism of Banerji and Gangulee (1937), Dharurkar (1971 Ph.D. Thesis), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis)- A Critical Review. *Environ. Conservation J.*, 8: 9-16.

- Salgare, S.A., 2007h. Alteration of resting period of pollen of five cultivars of the Apocynaceae by herbicide (2,4-Dinitrophenol): Further Evidence of a Criticism of Brewbaker and Kwack (1963), Sudhakaran (1967-Ph.D.Thesis), Saoji and Chitaley (1972), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review\*. Res. Hunt, 2: 107-114.
- Salgare, S.A., 2007i. Whether Optimum Pollen Germination and Tube Length Attained in the same growth medium (sucrose + sodium penta chloro phenate) by five cultivars of Apocynaceae [Further Evidence of a Criticism of Banerji and Gangulee (1937), Sudhakaran (1967-Ph.D.Thesis), Dharurkar (1971 Ph.D. Thesis), Nair, Nambudiri and Thomas (1973), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil and Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis)]. *Plant Archives*, 7: 403-408.
- Salgare, S.A., 2007j. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose+ basalin EC) by five cultivars of *Petunia axillaris* BSP.: Further evidence of a criticism of Banerji and Gangulee (1937), Sudhakaran (1967-Ph.D.Thesis), Dharurkar (1971 Ph.D. Thesis), Nair, Nambudiri and Thomas (1973), Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review. *Environ. Conservation J.*, 8: 21-30.
- Salgare, S.A., 2007k. Effect of fernoxone on the rate of pollen germination and tube growth of successive flowers of five cultivars of Petunia axillaris BSP.: Further Evidence of a Criticism of Saoji and Chitaley (1972), Berg (1973), Brandt (1974), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D.Thesis) and Giridhar (1984-Ph.D. Thesis) A Critical Review. Internat. J. Plant Sci., 2: 231-234.
- Salgare, S.A., 2007l. Monitoring of Herbicide (atrataf 50W) Toxicity by using Pollen as Indicators- Pollen of five cultivars of *Petunia axillaris* BSP.: Further Evidence of a Criticism of Berg (1973), Brandt



- (1974), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D.Thesis) and Giridhar (1984-Ph.D. Thesis)- A Critical Review. *Asian J. Bio Sci.*, 1: 191-195.
- Salgare, S.A. and Bindu, 2002. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose+ calcium chloride) by pinkflowered *Catharanthus roseus*: A Criticism to the Hypothesis of Nair, Nambudiri and Thomas (1973)— A Critical Review- I. *Biojournal*, 14: 21-23.
- Salgare, S.A., and Bindu, 2005. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose + calcium chloride) by white - flowered Catharanthus roseus and Further Evidence of A Criticism on the Hypothesis of Nair, Nambudiri and Thomas (1973)- A Critical Review. Flora and Fauna, 11: 183-184.
- Salgare, S.A. and Phunguskar, K.P., 2000. Monitors of pesticide (alphamethrin) toxicity by using pollen as indicators- Pollen of pink-flowered *Catharanthus roseus* A Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan, (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldara (1980), Shetye (1982- Ph.D. Thesis) and Giridhar (1984- Ph.D. Thesis) A Critical Review- III. *Biojournal*, 14: 1-4.
- Salgare, S.A. and Sanchita Pathak, 2005. Monitoring of heavy metal (bismuth nitrate) toxicity by using pollen as indicators Pollen of white-flowered *Catharanthus roseus* and further evidence of a Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis), and Giridhar (1984 Ph.D. Thesis) A Critical Review. *Flora and Fauna*, 11: 69-70.
- Salgare, S.A. and Sanju Singh, 2002. Monitors of heavy metal (ferrous sulphate) toxicity by using pollen as indicators- Pollen of white-flowered Catharanthus roseus- A Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldara (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis) A Critical Review Il. Biojournal, 14: 5-7.

- Salgare, S.A. and Sanju, Singh, 2006a. Monitoring of heavy metal (copper sulphate) toxicity by using pollen as indicators- Pollen of white-flowered Catharanthus roseus and further evidence of a Criticism on the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982- Ph.D. Thesis), and Giridhar (1984-Ph.D. Thesis)- A Critical Review- I. Him. J. Env. Zool., 20: 276-278.
- Salgare, S.A. and Sanju Singh, 2006b. Monitoring of heavy metal (cobalt nitrate) toxicity by using pollen as indicators- Pollen of pink-flowered cultivar of Catharanthus roseus and further evidence of a Criticism of the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980-Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982-Ph.D. Thesis) and Giridhar (1984-Ph.D. Thesis)- A Critical Review-II. Him. J. Env. Zool., 20: 283-284.
- Salgare, S.A. and Tessy, Mol Antony, 2005a. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose+ potassium chloride) by pink-flowered *Catharanthus roseus* and Further Evidence of A Criticism to the Hypothesis of Nair, Nambudiri and Thomas (1973)- A Critical Review. *Flora and Fauna*, 11: 61-62.
- Salgare, S.A. and Tessy Mol Antony, 2005b. Whether optimum pollen germination and tube length attained in the same growth medium (sucrose+ sodium oxalate) by white-flowered Catharanthus roseus and Further Evidence of A Criticism to the Hypothesis of Nair, Nambudiri and Thomas (1973)- A Critical Review. Flora and Fauna, 11: 75-76.
- Salgare, S.A. and Theresa, Sebastian, 1986. Further evidence of a Criticism of the hypothesis of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath and Kaleta (1978), Mhatre (1980 Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 Ph.D. Thesis) and Giridhar (1984 Ph.D. Thesis). 1st Nat. Symp. on Environ. Biol., held on December 30-31, 1986 at Deptt. Zool. and Microbiol., S. K. Univ., Anantapur- 515 003. Abstract No. 15.
- Salgare, S.A. and Theresa, Sebastian, 2006. Monitoring of herbicide (dalapon) toxicity by using pollen as indicators - Pollen of urid and further evidence of a Criticism of Berg (1973), Brandt (1974), Vick and Bevan (1976), Rasmussan (1977), Navara, Horvath



and Kaleta (1978), Mhatre (1980 - Ph.D. Thesis), Mhatre, Chaphekar, Ramani Rao, Patil, Haldar (1980), Shetye (1982 - Ph.D. Thesis) and Giridhar (1984 - Ph.D.Thesis) - A Critical Review - I\*. *Internat. J. Biosci. Reporter*, 4: 323-329.

Sharma, R.I., 1984. Effect of herbicides on plants of the Solanaceae - II. Ph.D. Thesis, Univ. Bombay.

- Shetye, R.P., 1982. Effect of heavy metals on plants. Ph.D. Thesis Univ. Bombay.
- Sudhakaran, I.V., 1967. Cytogenetic studies in Vinca rosea Linn. Ph.D. Thesis, Univ. Bombay.
- Trisa Palathingal, 1990. Evaluation of industrial pollution of Bombay by pollen- I. M. Phil. Thesis, Univ. Mumbai.

